

History

- Age
- Time of last meal
- Last bowel movement/ emesis
- Improvement or worsening with food or activity
- Duration of problem
- Other sick contacts
- **SAMPLE**
- Past surgical history
- Medications
- Menstrual history (pregnancy)
- Travel history
- Bloody emesis / diarrhea

Signs and Symptoms

- Pain
- Character of pain (constant, intermittent, sharp, dull, etc.)
- Distention
- Constipation
- Diarrhea
- Anorexia
- Radiation


Associated symptoms:

(Helpful to localize source)

Fever, headache, blurred vision, weakness, malaise, myalgias, cough, headache, dysuria, mental status changes, rash

Differential

- CNS (increased pressure, headache, stroke, CNS lesions, trauma or hemorrhage, vestibular)
- Myocardial infarction
- Drugs (NSAID's, antibiotics, narcotics, chemotherapy)
- GI or Renal disorders
- Diabetic ketoacidosis
- Gynecologic disease (ovarian cyst, PID)
- Infections (pneumonia, influenza)
- Electrolyte abnormalities
- Food or toxin induced
- Medication or Substance abuse
- Pregnancy
- Psychological



Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- Choose the lower dose of Promethazine (Phenergan) for patients likely to experience complications from sedative effects (e.g., elderly, debilitated, etc.)
- Document the mental status and vital signs prior to administration of Promethazine (Phenergan).
- Beware of vomiting only in children. Pyloric stenosis, bowel obstruction, and CNS processes (bleeding, tumors, or increased CSF pressures) all often present with vomiting.